

Renormalization group approach to multiparticle density fluctuations

S. Hegyi

Research Institute for Particle and Nuclear Physics of the Hungarian Academy of Sciences,

H-1525 Budapest 114, P.O. Box 49. Hungary

(February 1, 2008)

Abstract

An iterative procedure is developed with the aim of constructing homogeneity rules for the distribution $P(\rho, \delta)$ of the particle density ρ at resolution scale δ . A single iteration step consists of a change in the normalization point of $P(\rho, \delta)$ followed by a rescaling. Similar transformation rule is introduced for density fluctuations contaminated by Poisson noise. Application of the iterative procedure is given for the Ginzburg-Landau description of phase-transition from the quark-gluon plasma and for random cascading models.

PACS numbers: 13.85.Hd, 11.10.Gh, 05.40.+j

Over the past ten years significant changes have occurred in our knowledge of how to analyse the multiparticle density fluctuations and correlations, see Ref. [1] for a recent review. Several new techniques have been developed dominantly with the same goal: to search for self-similar patterns possessing no characteristic scale over a certain range of resolving power. Such fluctuations are expected to occur e.g. in elementary collision processes due to scale-invariant branching of parton cascades or in heavy-ion collisions as remnants of the phase transition from quark-gluon plasma to hadronic matter.

Self-similarity manifests itself in the power-law dependence of quantities characterizing the strength of fluctuations or correlations as the resolution scale (bin size in phase-space) is varied [1]. Besides power laws, there is another possible source of scaling behavior in multiparticle production. The probability distribution $P(\rho, \delta)$ of the particle density ρ at resolution scale δ can be a generalized homogeneous function of its arguments, $P(\rho, \delta) \sim \delta^{-\beta} \mathcal{F}(u = \rho/\delta^\nu)$ with β and ν being certain parameters [1]. In this case scaling refers to the data collapsing feature of homogeneous functions: after suitable rescaling the data points corresponding to different distributions $P(\rho, \delta)$ collapse onto a single curve, $\mathcal{F}(u)$.

Homogeneity rules play a central role in the theory of critical phenomena. Near the critical point of a physical system the thermodynamic functions exhibit homogeneous form [2] which implies the existence of a law of corresponding states. Using a suitably chosen scaling transformation it is possible to bring different states of the same system to coincidence and thus to compress many experimental or theoretical results into a compact form. This scaling hypothesis was first justified by the spin decimation procedure [3], the prototype of renormalization group (RG) transformations in statistical physics [4,5].

The connection between scale-invariant density fluctuations in multiparticle dynamics and the scaling laws of statistical physics has already been established [6,7]. In this Letter an attempt is made to deepen the relationship between the two fields by developing an iterative procedure that acts on the distribution $P(\rho, \delta)$ and provides the analogue of RG transformations. In the course of the iteration one may (and will) arrive at scaling behavior of $P(\rho, \delta)$ even if the distribution is not a homogeneous function of its arguments at the

beginning of the procedure.

Let us start with some elementary properties of the probability distribution $P(\rho, \delta)$ of observing particle density ρ at resolution scale (bin size) δ . For simplicity we shall consider here only averaging over the events. The effect of statistical noise will be discussed later on. The probability density $P(\rho, \delta)$ fulfills the normalization condition $\int_0^\infty P(\rho, \delta) d\rho = 1$. Its moments are defined by $\langle \rho^q(\delta) \rangle = \int_0^\infty \rho^q P(\rho, \delta) d\rho$. If only the first moment $\langle \rho(\delta) \rangle$ is responsible for the δ -dependence of $P(\rho, \delta)$ the distribution can be written in the homogeneous form

$$P(\rho, \delta) = \frac{1}{\langle \rho(\delta) \rangle} \mathcal{F}\left(\frac{\rho}{\langle \rho(\delta) \rangle}\right) \quad (1)$$

where the function $\mathcal{F}$ depends only on the scaling variable $u = \rho/\langle \rho(\delta) \rangle$. It satisfies the two normalization conditions $\int_0^\infty \mathcal{F}(u) du = \int_0^\infty u \mathcal{F}(u) du = 1$. Thus we have two properly normalized scaling functions, the second one is $\tilde{\mathcal{F}}(u) = u \mathcal{F}(u) = \rho P(u)$. Obviously, the normalized moments of $P(\rho, \delta)$ coincide with the moments of $\mathcal{F}(u)$,

$$\frac{\langle \rho^q(\delta) \rangle}{\langle \rho(\delta) \rangle^q} = \int_0^\infty u^q \mathcal{F}(u) du = C_q \quad (2)$$

which are independent of δ . Moreover, the moments of $\tilde{\mathcal{F}}(u)$ are given by $\tilde{C}_q = \int_0^\infty u^q \tilde{\mathcal{F}}(u) du = C_{q+1}$ i.e. the difference between the two scaling functions amounts to a shift in the ranks of their moments.

Although the scaling law of type Eq. (1) is frequently encountered in nature, for self-similar multiparticle density fluctuations it can not be valid. The observations indicate power-law dependence of the normalized moments of $P(\rho, \delta)$ as the resolution scale is varied, $C_q(\delta) \propto \delta^{-\varphi_q}$ where the φ_q are the intermittency exponents [1]. Accordingly, we have δ -dependent “scaling” functions $\mathcal{F}(u, \delta)$ and $\tilde{\mathcal{F}}(u, \delta)$. Let us observe that the latter enables one to perform a rescaling of type Eq. (1) yet again, the new scaling hypothesis reads

$$\tilde{\mathcal{F}}(u, \delta) = \frac{1}{\tilde{C}_1(\delta)} \mathcal{F}_1\left(\frac{u}{\tilde{C}_1(\delta)}\right). \quad (3)$$

(Rescaling $\mathcal{F}(u, \delta)$ in a similar manner has no effect since $C_1(\delta) = 1$.) The subscript of the new scaling function refers to the first step of an iterative sequence of transformations acting

on the distribution $P(\rho, \delta)$. For the starting distributions and moments corresponding to the 0th step we shall not indicate this explicitly.

Our new scaling variable is $u_1 = u/\tilde{C}_1(\delta)$ and the first-iterate moments $C_{q,1} = \int_0^\infty u_1^q \mathcal{F}_1(u_1) du_1$ are related to the original ones through

$$C_{q,1} = \frac{\tilde{C}_q(\delta)}{[\tilde{C}_1(\delta)]^q} = \frac{C_{q+1}(\delta)}{[C_2(\delta)]^q}. \quad (4)$$

Expressing now the scaling function $\mathcal{F}_1(u_1)$ similarly we get $\mathcal{F}_1(u_1) = C_2(\delta) \rho P[u/C_2(\delta)]$ which can be rewritten as

$$\mathcal{F}_1(u_1) = \frac{\langle \rho^2(\delta) \rangle}{\langle \rho(\delta) \rangle} \frac{\rho}{\langle \rho(\delta) \rangle} P\left(\rho \left/ \frac{\langle \rho^2(\delta) \rangle}{\langle \rho(\delta) \rangle} \right.\right). \quad (5)$$

It is seen that $\mathcal{F}_1(u_1)$ has the same form as $\mathcal{F}(u)$, namely, $\mathcal{F}_1(u_1) = \langle \rho(\delta) \rangle_1 P_1[\rho/\langle \rho(\delta) \rangle_1]$ where the first-iterate distribution $P_1(\rho, \delta)$ and its moments $\langle \rho^q(\delta) \rangle_1$ are obtained by changing the normalization point of $P(\rho, \delta)$ from the 0th moment to $\langle \rho(\delta) \rangle$ and rescaling by $\langle \rho(\delta) \rangle$ to maintain the overall normalization.

Demonstrating the benefits of the above transformation of $P(\rho, \delta)$ let us assume that the self-similar density fluctuations show monofractal properties. Then the intermittency exponents are given by $\varphi_q = \varphi_2(q-1)$ and the anomalous fractal dimensions $D_q = \varphi_q/(q-1)$ are q -independent, $D_q = D_2$. The normalized moments of $P(\rho, \delta)$ take the form $C_q(\delta) = A_q[C_2(\delta)]^{q-1}$ with coefficients A_q independent of δ . However, Eq. (4) yields $A_q = C_{q-1,1}$. This implies that monofractal density fluctuations give rise not only to power-law scaling of the normalized moments of $P(\rho, \delta)$ but also to homogeneous form of the distribution after changing its normalization point to $\langle \rho(\delta) \rangle$ and making an appropriate rescaling.

In the followings we omit explicit reference to the variable δ to simplify the notation. If $\mathcal{F}_1(u_1)$ is not independent of the resolution scale, the single iteration step described before can be repeated until the appearance of scaling (i.e. data collapsing) behavior provided that certain moments of $P(\rho)$ exist, see below. In the k th step of the iteration we have the normalization conditions $\int_0^\infty \mathcal{F}_k(u_k) du_k = \int_0^\infty \tilde{\mathcal{F}}_k(u_k) du_k = 1$ where $\tilde{\mathcal{F}}_k(u_k) = u_k \mathcal{F}_k(u_k)$. The distribution $\mathcal{F}_k(u_k) = \langle \rho \rangle_k P_k(\rho/\langle \rho \rangle_k)$ is related to the $(k-1)$ th step through

$$\mathcal{F}_k(u_k) = \tilde{C}_{1,k-1} \tilde{\mathcal{F}}_{k-1} \left(\frac{u_{k-1}}{\tilde{C}_{1,k-1}} \right) \quad (6)$$

and its moments $C_{q,k} = \int_0^\infty u_k^q \mathcal{F}_k(u_k) du_k$ are given by

$$C_{q,k} = \frac{\tilde{C}_{q,k-1}}{\tilde{C}_{1,k-1}^q} = \frac{C_{q+1,k-1}}{C_{2,k-1}^q}. \quad (7)$$

The $(k+1)$ th step of the iteration follows in the same way as the first step in Eq. (3),

$$\tilde{\mathcal{F}}_k(u_k) = \frac{1}{\tilde{C}_{1,k}} \mathcal{F}_{k+1} \left(\frac{u_k}{\tilde{C}_{1,k}} \right). \quad (8)$$

Scaling behavior of $P_{k+1}(\rho)$ appears if the monofractal type relationship $C_{q,k} \propto C_{2,k}^{q-1}$ holds with constants of proportionality independent of the resolution scale δ .

Let us now consider how the distribution $P_k(\rho)$ is related to the original $P(\rho)$. The connection is provided by the straightforward extension of the normalization point changing transformation discussed previously,

$$P_k(\rho) = \frac{\rho^k}{\langle \rho^k \rangle} P(\rho) \quad (9)$$

and thus the moments of $P_k(\rho)$ are given by $\langle \rho^q \rangle_k = \langle \rho^{q+k} \rangle / \langle \rho^k \rangle$. From this we get for the normalized moments $C_{q,k}$ the expression

$$C_{q,k} = \frac{\langle \rho^q \rangle_k}{\langle \rho \rangle_k^q} = C_{q+k} \frac{C_k^{q-1}}{C_{k+1}^q} \quad (10)$$

in terms of the original normalized moments C_q . Homogeneity of $P_{k+1}(\rho)$ appears in case of δ -independence of the product $C_{k+q} C_{k+2}^{1-q} C_{k+1}^{q-2}$. The necessary condition of performing the k th iteration step is the existence of the moments of $P(\rho)$ up to k th order.

Eq. (9) is analogous to the RG transformations which correspond in many important cases to a change in the norm of the parameters characterizing a physical system [5]. It is a Gell-Mann and Low type relationship [8] in which a multiplicative transformation (in our case multiplication by ρ^k) is compensated by a rescaling and a suitable change in the renormalized parameters. Since the operation $P_k(\rho) \rightarrow P_{k-1}(\rho)$ does not exist the iteration steps constitute a semi-group. In statistical physics the Kadanoff-Wilson type RG transformations

perform a systematic reduction in the number of degrees of freedom (spin decimation). This enables one to eliminate the small-scale fluctuations from the problem which are irrelevant to critical point behavior such as the homogeneity of thermodynamic functions. In the iterative procedure developed here the elimination of fluctuations manifests through elimination of the moments of $P(\rho)$ via moment-shifting and rescaling until one observes a homogeneity rule of type Eq. (1) for $P_k(\rho)$.

The finite number of detected particles gives rise to statistical fluctuations contaminating the distribution $P(\rho)$ of purely dynamical origin. It is convenient to represent the statistical component by a Poisson process whose convolution with $P(\rho)$ can be described by the Poisson transform [1]. It yields the discrete probability distribution $P(n)$ of detecting n particles:

$$P(n) = \int_0^\infty P(\rho) \frac{\rho^n}{n!} e^{-\rho} d\rho. \quad (11)$$

The Poisson transform has the useful property that the factorial moments ξ_q of $P(n)$ are equivalent to the ordinary moments of $P(\rho)$,

$$\xi_q = \langle n^{[q]} \rangle = \sum_{n=q}^\infty n^{[q]} P(n) = \langle \rho^q \rangle \quad (12)$$

where $n^{[q]} = n(n-1)\dots(n-q+1)$. Thus in the presence of Poisson noise factorial observables make it possible to study directly the dynamical fluctuations whose self-similarity is signalled by the power-law dependence of the normalized factorial moments $F_q = \xi_q/\xi_1^q$ on the resolution scale δ [9].

We are interested here in the Poisson transform of the iterated dynamical distribution $P_k(\rho)$. But let us first consider briefly some properties of Eq. (9) itself. This transformation of $P(\rho)$ is known as size-biasing of order k in the mathematical literature [10]. A worth noting feature is that the log-exponential family of continuous probability densities are form-invariant under size-biasing, only their parameters are affected. In this respect the log-normal distribution deserves special attention. Its parameters are changed by size-biasing of order k according to $\mu \rightarrow \mu + k\sigma^2$ with μ and σ^2 being the mean and variance of the

original distribution. Therefore a log-normal $P(\rho)$ is a fixed point of our iterative procedure: although Eq. (9) changes its first moment, this change is scaled out by constructing $\mathcal{F}_k(u_k)$ and we get an unaffected scaling function at each iteration step.

The Poisson transform of $P_k(\rho)$ leads to size-biasing of factorial order k acting on $P(n)$:

$$P_k(n) = \int_0^\infty P_k(\rho) \frac{\rho^n}{n!} e^{-\rho} d\rho = \frac{n^{[k]}}{\langle n^{[k]} \rangle} P(n) \Big|_{\Leftarrow k} \quad (13)$$

where the symbol $\Leftarrow k$ denotes left-shifting the size-biased $P(n)$ by k places. It is seen that we have the same kind of transformation as for $P(\rho)$: the normalization point of $P(n)$ is changed to $\langle n^{[k]} \rangle$ and a rescaling by $\langle n^{[k]} \rangle$ is performed to maintain the overall normalization. This yields a discrete distribution with k leading zeroes, $P(n < k) = 0$, hence a left-shift by k places is required to maintain also the original support in n of the nonzero probabilities [10]. The normalized factorial moments $F_{q,k}$ are related to F_q in the same way as $C_{q,k}$ to C_q i.e. $F_{q,k} = F_{q+k} F_k^{q-1} F_{k+1}^{-q}$. The expression for $P_k(n)$ implies that the normalization point changing transformations form a discrete semi-group in the presence of Poisson noise. On the contrary, $P_k(\rho)$ is well defined for fractional k too although the iterative procedure that led us to Eq. (9) enables only integer valued k . It is worth mentioning that $F_{2,k-2} = F_k F_{k-2} F_{k-1}^{-2}$ yields for $k \geq 2$ the k th bunching parameter [11] in the limit $\delta \rightarrow 0$.

We now turn our attention to applications of our results. In recent years considerable interest has been devoted to the Ginzburg-Landau theory of the phase transition from quark-gluon plasma to hadronic matter [12,13]. Instead of a strict power-law scaling $F_q \propto \delta^{-\varphi_q}$ of the normalized factorial moments, a so-called F -scaling behavior $F_q \propto F_2^{\beta_q}$ is observed with $\beta_q = (q-1)^\nu$. For second-order transition $\nu = 1.304$ and in the case of a first-order transition $\nu = 1.45$. The F -scaling rule of the above form has the property $F_0 \neq 1$ hence it is difficult to identify the family of discrete probability laws that belongs to the observed F_q -pattern. Performing a single iteration step the factorial moments change according to $F_{q,1} \propto F_2^{\beta_q}$ with $\beta_q = q^\nu - q$ and therefore

$$F_{q,1} \propto F_{2,1}^{\beta_q} \quad \text{with} \quad \beta_q = \frac{q^\nu - q}{2^\nu - 2}. \quad (14)$$

This form of F -scaling is the log-Lévy law [15] with Lévy index $0 < \nu \leq 2$. We have thus identified the family of probability laws for $P_1(n)$ in the Ginzburg-Landau formalism with a strict bound on ν . The theoretically relevant values are within the allowed range. Our result can be checked in a laser experiment at the threshold of lasing where the photocount fluctuations are in accordance with the Ginzburg-Landau theory of second-order phase transitions and the F -scaling rule with $\nu = 1.304$ has been experimentally confirmed [14].

The log-Lévy law $F_q \propto F_2^{\beta_q}$ with β_q given in Eq. (14) is introduced to describe multiplicative random cascading models [15]. Let us keep track of the evolution of factorial moments in the course of successive iteration steps. It can be characterized by the ratio

$$\frac{D_{q,k}}{D_q} = \frac{(q+k)^\nu + (q-1)k^\nu - q(k+1)^\nu}{q^\nu - q} \quad (15)$$

whose k -dependence is shown in Fig. 1 for $q = 2$ and for different values of the Lévy index ν . Smaller is the value of ν faster is the convergence to $D_{2,k} = 0$, i.e. to δ -independence of $F_{2,k}$. This is what we expect since $\nu = 0$ corresponds to monofractal behavior which yields δ -independent factorial moments after a single iteration step. The $\nu = 2$ limiting case of the Lévy index defines maximum degree of multifractality. Random cascades of this type are described by the log-normal law whose fixed point property discussed before is apparent from Fig. 1. For $\nu > 2$, which is not permitted for log-Lévy laws, $F_{2,k}$ gets farther away from δ -independence.

Our last example concerns the analogy between fractal structures and statistical mechanics [1,16]. In the thermodynamic description of multifractal cascades the rank q plays the role of inverse temperature [16]. Analogously to RG calculations in statistical physics, performing the iterative sequence of transformations can be thought as varying the temperature. The evolution of $D_{q,k}$ allows the introduction of an effective inverse temperature q_k whose variation with k can be obtained from the equation $(q+k)^\nu + (q-1)k^\nu - q(k+1)^\nu = q_k^\nu - q_k$. The Lévy index ν controls the rate at which the effective inverse temperature changes in the course of the iteration. Therefore ν plays the role of the critical exponent of the correlation length characterizing condensed matter systems at the critical temperature [5].

In conclusion, we have developed an iterative procedure capable of finding homogeneity rules for a distribution that describes the fluctuation properties of a certain physical system. A single iteration step consists of a change in the normalization point of the distribution compensated by a suitable change in the renormalized parameters and a rescaling. The number of steps required to arrive at data collapsing behavior measures how strongly the scaling hypothesis of type Eq. (1) is violated. For self-similar fluctuations it is maximal in the case of maximum degree of multifractality, providing the fixed point of the iteration, and minimal for monofractals which give rise to scaling behavior after a single iteration step. Concerning multiparticle dynamics, our results shed new light on how to characterize the degree of KNO scaling violation of the full phase-space multiplicity distributions [17] in terms of novel scaling laws. Since homogeneity rules and data collapsing deserve special attention in many areas of physics the iterative procedure can have a number of applications in other fields as well.

This research was supported by the Hungarian Science Foundation under grant OTKA-F4019/1992.

REFERENCES

- [1] E.A. De Wolf, I.M. Dremin and W. Kittel, Phys. Rep. **270**, 1 (1996).
- [2] B.J. Widom, J. Chem. Phys. **43**, 3892, 3898 (1965).
- [3] L.P. Kadanoff, Physics **2**, 263 (1966).
- [4] K.G. Wilson and J. Kogut, Phys. Rep. **12C**, 75 (1974).
- [5] S.K. Ma, *Modern Theory of Critical Phenomena* (Benjamin, Reading, MA, 1976).
- [6] R.C. Hwa, in *Quark-Gluon Plasma*, edited by R.C. Hwa (World Scientific, Singapore, 1990).
- [7] J. Wosiek, in *Proc. XXIV Int. Symp. on Multiparticle Dynamics*, edited by A. Giovannini, S. Lupia and R. Ugoccioni (World Scientific, Singapore, 1995).
- [8] M. Gell-Mann and F.E. Low, Phys. Rev. **95**, 1300 (1954).
- [9] A. Białas and R. Peschanski, Nucl. Phys. **B273**, 703 (1986); **B308**, 857 (1988).
- [10] G.P. Patil and J.K. Ord, Sankhyā **B38**, 48 (1976).
- [11] S.V. Chekanov and V.L. Kuvshinov, Acta Phys. Pol. **B25**, 1189 (1994).
- [12] R.C. Hwa and M.T. Nazirov, Phys. Rev. Lett. **69**, 741 (1992).
- [13] R.C. Hwa, Phys. Rev. **D47**, 2773 (1993); Phys. Rev. **C50**, 383 (1994).
- [14] M.R. Young, Y. Qu, S. Singh and R.C. Hwa, Optics Commun. **105**, 325 (1994).
- [15] Ph. Brax and R. Peschanski, Phys. Lett. **B253**, 225 (1991).
- [16] R. Peschanski, Nucl. Phys. **B327**, 144 (1989).
- [17] Z. Koba, H.B. Nielsen and P. Olesen, Nucl. Phys. **B40**, 314 (1972).

FIGURES




FIG. 1. Variation of the anomalous fractal dimension D_2 with iteration step k for different values of the Lévy index ν .